


Journal Pre-proof

Research Note: A mixture of *Bacteroides* spp. and other probiotic intestinal anaerobes reduces colonization by pathogenic *E. coli* strain O78:H4-ST117 in newly hatched chickens

Aneta Papouskova , Ivan Rychlik , Danka Harustiakova ,
Alois Cizek

PII: S0032-5791(23)00053-6
DOI: <https://doi.org/10.1016/j.psj.2023.102529>
Reference: PSJ 102529



To appear in: *Poultry Science*

Received date: 22 November 2022
Accepted date: 18 January 2023

Please cite this article as: Aneta Papouskova , Ivan Rychlik , Danka Harustiakova , Alois Cizek , Research Note: A mixture of *Bacteroides* spp. and other probiotic intestinal anaerobes reduces colonization by pathogenic *E. coli* strain O78:H4-ST117 in newly hatched chickens, *Poultry Science* (2023), doi: <https://doi.org/10.1016/j.psj.2023.102529>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 Published by Elsevier Inc. on behalf of Poultry Science Association Inc.
This is an open access article under the CC BY-NC-ND license
(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Research Note: A mixture of *Bacteroides* spp. and other probiotic intestinal anaerobes reduces colonization by pathogenic *E. coli* strain O78:H4-ST117 in newly hatched chickens

Aneta Papouskova^{*,1}, Ivan Rychlik[#], Danka Harustiakova[§], Alois Cizek^{*}

*Institute of Microbiology and Infectious Diseases, Faculty of Veterinary Medicine, University of Veterinary Sciences Brno, Brno, The Czech Republic

[#]Veterinary Research Institute, Brno, The Czech Republic

[§] RECETOX, Faculty of Science, Masaryk University, Brno, The Czech Republic

¹ Corresponding author:

Aneta Papouskova, Institute of Microbiology and Infectious Diseases, Faculty of Veterinary Medicine, Veterinary University Brno, Brno, Czech Republic

email: papouskovaa@vfu.cz

telephone: +420 541562278

Conflict of interest statement: The authors declare no conflict of interest.

Summary

An experimental group of one-day-old chicken from a commercial hatchery was given a defined mixture of 7 gut anaerobes. The next day the chicks were inoculated by an APEC strain O78:H4-ST117 resistant to ciprofloxacin, alongside with the control group and

monitored for 4 weeks after the inoculation for the presence of the colonizing strains and ciprofloxacin-resistant *E. coli*. Significant reduction of colonization rates in the first two weeks was recorded in the experimental group for the numbers of ciprofloxacin-resistant *E. coli*. The results show that colonization of chicken by defined anaerobic mixtures may provide a decisive protection during the critical period of the chicken intestinal microflora development.

Keywords: competitive exclusion; colonization; chicken; avian pathogenic *Escherichia coli* (APEC); probiotics

INTRODUCTION

The importance of colibacillosis as a paramount economic and public health issue is well recognized. *E. coli* strains that are associated with colibacillosis have been termed APEC, avian pathogenic *E. coli*, and usually belong to certain serotypes and are characteristic by the presence of specific virulence genes. APEC are thought to form a specific pathotype within the group of extraintestinal pathogenic *E. coli* (ExPEC). Unfortunately, the immense diversity of these strains, many of which lack typical traits, complicate the precise definition of the pathotype. Recent studies confirm the importance of both secondary infections caused by very diverse commensal-like strains and the occurrence of adapted virulent lineages, often with presumable zoonotic potential (Mehat et al., 2021). In addition, virulent lineages may spread vertically through the production pyramid from breeder farms to production flocks via hatcheries (Projahn et al., 2017).

Intervention measures therefore pose a challenge. Use of antibiotics only supports the emergence and spread of resistant strains. In addition, the pressure to limit or completely ban antimicrobials in food animals may result in substantial losses in production farms and, on the other hand, calls for a discussion on different, more sustainable means of production. The principle of competitive exclusion (CE) represents one of the most promising strategies for reducing infectious pressure and transmission of pathogenic strains. Since its introduction in 1973 (Rantala et Nurmi, 1973), commercial products such as Aviguard and Broilact have been developed and their efficacy confirmed not only against *Salmonella* sp., but also against *E. coli* producing extended-spectrum beta-lactamases (ESBL) (Methner et al., 2019) and pathogenic *E. coli* in broilers (Hofacre et al., 2002).

Induction of "mature" intestinal microflora facilitates the development of the intestinal mucosal immune system and increases resistance to *Salmonella* infection (Crhanova et al., 2011). Notably, complex, often only partially defined, mixtures of intestinal anaerobic bacteria or cecal transplants have proven to be much more effective than a single species probiotic or a limited combination of several traditional probiotic strains, most of which are incapable of long-term colonization. Traditional probiotics mainly include lactobacilli, but their effects are often questionable (Yan et al., 2017; Li et al., 2021). The application of strict anaerobes, mainly from the phylum Bacteroidetes therefore makes more sense than administering facultative anaerobes and spore-forming anaerobes of which the environment is an abundant source (Karasova et al., 2022; Kubasova et al., 2019a; Kubasova et al., 2021).

The purpose of this study was to evaluate the effect of a complex mixture of defined intestinal anaerobes on cecal colonization by ciprofloxacin-resistant strain APEC ST117-O78:H4, a representative of one of the most prevalent risk lineages in poultry industry (Ronco et al., 2017).

MATERIAL AND METHODS

Experimental animals

Ross 308 meat hybrid chickens were used for the experiment. The one-day-old chickens were delivered from a local commercial hatchery, originated from one parent farm and were formed by a mixture of both sexes.

Housing conditions and care

The experimental and control groups, each consisting of 28 chicks, were placed in two separate rooms on sawdust bedding, near the heating lamp. The commercial feed for given age was provided *ad libitum*. Water with the addition of a vitamins and organic acids (Acidomide, dosage according to the manufacturer's recommendations) was also administered *ad libitum*. A 12-hour light regime was followed and the ambient temperature was recorded regularly.

Probiotic mixture and colonization *E. coli* strain

The probiotic mixture consisted of 7 bacterial species including *Bacteroides caecicola*, *Bacteroides plebeius*, *Megasphaera stantonii*, *Megamonas hypermegale*, *Megamonas funiformis*, *Phascolarctobacterium faecium* and *Sutterella massiliensis*. These strains were inoculated into 100 ml of Wilkins-Chalgren nutrient broth and incubated in anaerobic cabinet at 37 °C in atmosphere consisting of 85 % N₂, 10 % CO₂ and 5 % H₂. After 48-hour-fermentation, 0.1 ml of complex bacterial culture was used for oral inoculation of each chick

in experimental group. An aliquot of 1 ml of the complex bacterial culture was saved and used for DNA purification to determine ratios of each of the strains in the inoculum.

E. coli 587 (serotype O78:H4, ST117) was used for challenge. This strain, representing an important APEC line, had been analyzed by whole-genome sequencing previously (Papouskova et al., 2020). The experimental strain was cultured in BHI at 37 °C for 18 hours and the culture was diluted with sterile buffered saline to a concentration of 10^5 CFU/ml as verified by cultivation. Each chicken was gavaged with 0.1 ml of inoculum, i.e. with a dose of 10^4 CFU.

The course of the experiment

The experiment lasted four weeks. On day 1 of life, chicks in the experimental group were orally inoculated with 0.1 ml of gut anaerobes. On the day 2, both groups received a suspension of ciprofloxacin-resistant strain *E. coli* 587. On days 7, 14, 21 and 28 of life, 7 chickens were randomly selected from each group, weighed and euthanized by carbon dioxide inhalation. Pathomorphological changes were recorded at the necropsy and the ceca of all chickens were aseptically extirped.

Verification of colonization rate by probiotic mixture

Samples of the administered probiotic mixture and caecal contents were stored at -20 °C. DNA from caecal samples was extracted using QIAamp Stool kit according to the manufacturer's instructions (Qiagen, Germany). Based on known genomic sequences, strain-specific real-time PCRs were designed and the real-time PCR in SybrGreen format was performed exactly as described previously (Kollarcikova et al, 2020).

Verification of colonization rate by challenge APEC strain

A serial decimal dilutions of cecal contents with sterile buffered saline were plated on Levine EMB agar (Thermo Fischer Diagnostics, Dardilly, France) supplemented with 1 µg/ml of ciprofloxacin (Sigma Aldrich, Prague, Czech Republic). After 18 hours of incubation at 37 °C, colonies with a typical metallic sheen were counted and the presence of *E. coli* O78 was verified by slide agglutination with specific antisera according to the manufacturer's recommendations (Denka-Seiken Co., Ltd., Japan). The presence of the experimental strain was verified by slide agglutination on at least two plates with sufficient growth, and according to the highest dilution at which strain O78 was still detected, a colonization score of + (dilutions 10^{-1} to 10^{-2}), ++ (dilutions 10^{-3} to 10^{-4}) and +++ (dilutions 10^{-5} to 10^{-6} , i.e. the highest), was assigned for easier result presentation.

Statistical analysis

The obtained experimental data (cfu.g⁻¹) were log₁₀ transformed, and the mean values and standard deviations were calculated. Factorial ANOVA followed by Tuckey post hoc test was used to assess the effect of control/experimental group and chicken age on number of bacteria in the gut. $P < 0.05$ was considered as significant in all tests. Statistical analyses were performed using Statistica, version 13 (TIBCO Software Inc.).

RESULTS AND DISCUSSION

The aim of the experiment was to verify the effect of application of the probiotic mixture to one-day-old chicks on their health, performance and resistance to colonization by a defined APEC strain. Probiotic mixture consisted of 7 different species, though because the strains

were co-fermented in a single culture medium, their representation was not identical. Instead, the inoculum was dominated by *Bacteroides caecicola*, *Bacteroides plebeius*, *Megasphaera stantonii* and *Sutterella massiliensis* while *Megamonas hypermegale*, *Megamonas funiformis* and *Phascolarctobacterium faecium* were present at lower than 0.2 % representation (Fig. 1A). As expected, all the strains efficiently colonised chicken caecum though the composition of the inoculum only partially affected caecal colonisation (Kubasova et al., 2019b).

Bacteroides caecicola and *Bacteroides plebeius* dominated also in the caecum but *Megasphaera stantonii* was less abundant in the caecum than in the inoculum (Fig. 1B).

Moreover, minority species from the inoculum formed more than 1 % of total microbiota in the caecum *in vivo*. Altogether, the used probiotic strains formed 48 % of total microbiota on average ranging from 32.4 to 55.9 % in individual chicks. All of the probiotic strains were absent from microbiota of control chicks (Fig. 1C).

No mortality was recorded during the experiment and the chickens showed no visible pathomorphological changes at necropsy. No effect of the probiotic mixture on the weight of the chickens was observed; the mean weight remained slightly higher in the experimental group, but this difference was not statistically significant (Table 1).

In the first two weeks, statistically significant differences in the numbers of all CIP-resistant *E. coli* between the control and experimental groups were recorded (Table 1). In the control group, the numbers of CIP-resistant *E. coli* reached values of 10^7 to 10^8 CFU/g, in the experimental group these were approximately two logs of magnitude lower. These data also show that the competitive exclusion effect was not specific against the pathogenic strain but was general, decreasing the colonisation of all *E. coli* clones present also in the control chickens. In older chickens (three-week and four-week-old), the numbers of *E. coli* in the gut

were significantly lower than in younger chickens, probably due to natural maturation and stabilization of the intestinal ecosystem, and these no longer differed between the control and experimental groups (Table 1).

The results of the quantitative cultivation of cecal contents were complicated by the presence of *E. coli* clones resistant to ciprofloxacin present also in the control chickens. The chickens must have been colonized by these strains either in the hatchery, during transport or shortly after placement in the animal house. CFU numbers varied by about two logs of magnitude in first two weeks, a result comparable to the effects of some single species probiotics, such as lactobacilli (Edens et al., 1997). Since the colonization by field *E. coli* strains probably occurred before the application of the probiotic mixture, a certain therapeutic effect was recorded, although further experimental work would be needed to confirm the higher efficacy of the probiotic mixture if applied directly in the hatchery. The results can also be compared with direct challenge with septicemic strain O78:H80 after application of a commercial competitive exclusion product (Aviguard), which led to a reduction in cecal colonization of 2-3 logs on days 7 and 14 (Hofacre et al., 2002) and studies testing the effect of the same product on ESBL/AmpC colonization producing *E. coli* strains (a decrease of 2-4 logs depending on the challenge strain dose) (Methner et al., 2019).

The experimental and control groups differed significantly in colonization by the experimental *E. coli*. The experimental strain was not detected in the experimental group when culturing the cecal contents with an exception of a single sample, which showed positive agglutination only at the lowest dilution; the experimental strain persisted in low intestinal numbers in this chicken at the week 4. In contrast, in the control group in the first two weeks, the experimental strain was found in all chickens up to a dilution of 10^5 to 10^6

CFU/g of cecal content (Tab. 1). In the third and fourth week, the O78 strain was recorded in about half of the chickens in the control group (in 3 and 4 chickens out of 7 tested, respectively). Thus, the probiotic mixture prevented colonization of the chickens in the experimental group by the APEC below a detectable level, whereas in the control group there was a massive and 100% colonization, which lasted for at least 14 days in all chickens. From the third week onwards, the differences between the groups were insignificant, but there was a significant decrease in *E. coli* colonization compared to previous weeks. The time course corresponds to the natural development of the intestinal microflora of commercially reared chickens, as previously described (Videnska et al., 2014), characterized by massive colonization by Proteobacteria in the first week and their gradual replacement by Firmicutes after the second week of life. The intestines of newly hatched chickens are susceptible to colonization by environmental bacteria, including potential pathogens, and the first hours of life determine the development of the intestinal ecosystem for next weeks. The first week is critical in terms of broiler chicken mortality, it is a period of extreme susceptibility to enteric and secondary systemic infections such as colibacillosis and salmonellosis (Crhanova et al., 2011). In contrast, chickens hatched in contact with adults very quickly develop "mature" intestinal microflora and associated resistance (Kubasova et al., 2019a). Although a time-limited effect can be expected after a single application of probiotics, colonization with complex microflora can be a way to ensure a substantial level of protection for commercial chickens in the most critical period. A satisfactory effect can be expected only i) with early application of the mixture, before colonization of the intestine by bacteria from the environment; administration on the fattening farm comes too late; ii) when using complex mixtures, i.e. commercial CE products, cecal content or own anaerobic mixtures, rather than classical probiotics such as lactobacilli, the effects of which are questionable.

Acknowledgements

Special thanks to Sona Krmickova and Vera Novakova for their help during experiments.

This experiment has been supported by projects QK22020066 of the Czech Ministry of Agriculture and CEITEC/Literak/ITA 2021 by the Veterinary University Brno. Authors thank to Research Infrastructure RECETOX RI (No LM2018121) financed by the Ministry of Education, Youth and Sports, and Operational Programme Research, Development and Innovation – project CETOCOEN EXCELLENCE (No CZ.02.1.01/0.0/0.0/17_043/0009632) for supportive background.

Conflict of interest statement

The authors declare no conflict of interest.

Ethic statement

The experiment was reviewed and approved by the committee for ethics in animal testing, Czech Ministry of Education, Youth and Sports, No. MSMT-19246/2021-3.

REFERENCES

- Crhanova, M., H. Hradecka, M. Faldynova, M. Matulova, H. Havlickova, F. Sisak, I. Rychlik. 2011. Immune response of chicken gut to natural colonization by gut microflora and to *Salmonella enterica* serovar enteritidis infection. *Infect Immun.* 79:2755-63.
- Edens F.W., C.R. Parkhurst, I.A. Casas, W.J. Dobrogosz. 1997. Principles of ex ovo competitive exclusion and in ovo administration of *Lactobacillus reuteri*. *Poult Sci.* 76:179-96.

- Hofacre C.L., A.C. Johnson, B.J. Kelly, R. Froyman. 2002. Effect of a commercial competitive exclusion culture on reduction of colonization of an antibiotic-resistant pathogenic *Escherichia coli* in day-old broiler chickens. *Avian Dis.* 46:198-202.
- Kollarčíková M., M. Faldynová, J. Matiasovicová, E. Jahodarová, T. Kubasová, Z. Seidlerová, V. Babák, P. Videnská, A. Cizek, I. Rychlík. 2020. Different *Bacteroides* Species Colonise Human and Chicken Intestinal Tract. *Microorganisms.* 8:1483.
- Karasová D., M. Faldynová, J. Matiasovicová, A. Sebková, M. Crhanová, T. Kubasová, Z. Seidlerová, H. Prikrylová, J. Volf, M. Zeman, V. Babák, H. Juricová, J. Rajová, L. Vlasatíková, P. Rysavka, I. Rychlík. 2022. Host species adaptation of obligate gut anaerobes is dependent on their environmental survival. *Microorganisms.* 10: 1085.
- Kubasová T., M. Kollarčíková, M. Crhanová, D. Karasová, D. Cejková, A. Sebková, J. Matiasovicová, M. Faldynová, A. Pokorná, A. Cizek, I. Rychlík. 2019a. Contact with adult hen affects development of caecal microbiota in newly hatched chicks. *PLoS One.* 14: e0212446.
- Kubasová T., M. Kollarčíková, M. Crhanová, D. Karasová, D. Cejková, A. Sebková, J. Matiasovicová, M. Faldynová, F. Sisák, V. Babák, A. Pokorná, A. Cizek, I. Rychlík. 2019b. Gut anaerobes capable of chicken caecum colonisation. *Microorganisms.* 7:597.
- Kubasová T., Z. Seidlerová, I. Rychlík. 2021. Ecological adaptations of gut microbiota members and their consequences for use as a new generation of probiotics. *Int J Mol Sci.* 22: 5471.
- Li T., C.D. Castañeda, J. Miotto, C. McDaniel, A.S. Kiess, L. Zhang. 2021. Effects of in ovo probiotic administration on the incidence of avian pathogenic *Escherichia coli* in broilers and an evaluation on its virulence and antimicrobial resistance properties. *Poult Sci.* 1003:100903.

- Mehat J.W., A.H.M. van Vliet, R.M. La Ragione. 2021. The Avian Pathogenic *Escherichia coli* (APEC) pathotype is comprised of multiple distinct, independent genotypes. *Avian Pathol.* 50:402-416.
- Methner U., A. Friese, U. Rösler. 2019. Competitive exclusion: A tool to combat extended-spectrum β -lactamase-producing *Escherichia coli* strains in chickens. *Res Vet Sci.* 123:124-128.
- Papouskova A., M. Masarikova, A. Valcek, D. Senk, D. Cejkova, E. Jahodarova, A. Cizek. 2020. Genomic analysis of *Escherichia coli* strains isolated from diseased chicken in the Czech Republic. *BMC Vet Res.* 16:189.
- Projahn M., K. Daehre, U. Roesler, A. Friese. 2017. Extended-Spectrum-Beta-Lactamase- and Plasmid-Encoded Cephamycinase-Producing Enterobacteria in the Broiler Hatchery as a Potential Mode of Pseudo-Vertical Transmission. *J Appl Environ Microbiol.* 83, e02364-16.
- Rantala M., E. Nurmi. 1973. Prevention of the growth of *Salmonella* Infantis in chicks by the flora of the alimentary tract of chickens. *Br Poult Sci.* 14:627-30.
- Ronco T., M. Stegger, R.H. Olsen, C. Sekse, A.B. Nordstoga, T. Pohjanvirta, B. Lilje, U. Lyhs, P.S. Andersen, K. Pedersen. 2017. Spread of avian pathogenic *Escherichia coli* ST117 O78:H4 in Nordic broiler production. *BMC Genomics.* 18:13.
- Videnska P., K. Sedlar, M. Lukac, M. Faldynova, L. Gerzova, D. Cejkova, F. Sisak, I. Rychlik. 2014. Succession and replacement of bacterial populations in the caecum of egg laying hens over their whole life. *PLoS One.* 9:e115142.
- Yan W., C. Sun, J. Yuan, N. Yang. 2017. Gut metagenomic analysis reveals prominent roles of *Lactobacillus* and cecal microbiota in chicken feed efficiency. *Sci Rep.* 7:45308.

Figure legend

Figure 1. Microbiota composition in the inoculum and caeca of inoculated and control chicks. Real time PCR was used to quantify probiotic strains in complex populations. The inoculum was dominated by *Bacteroides caecicola*, *Bacteroides plebeius*, *Megasphaera stantonii* and *Sutterella massiliensis* (Panel A). Ratios calculated only for the probiotic strains showed that composition of the inoculum only partially influenced caecal colonisation (Panel B). Used probiotic strains were completely absent from the caeca of control chicks (Panel C).


Table 1. Body weight of chickens and number of *E. coli* in the caecum of chickens in control (CTRL) and experimental (Exp) group.

	Body weight (g)	<i>E. coli</i> counts (log CFU/g of caecum) ^a	Experimental strain detection (number of positive chicken, colonization rate ^b)			
Week	Exp	CTRL	Exp	CTRL	Exp	CTRL
1	126.4 ± 8.2	111.0 ± 17.8	6.38 ± 0.45*	8.29 ± 0.70	ND	7/7 (+++)
2	306.4 ± 90.9	262.3 ± 61.9	6.04 ± 0.47*	7.83 ± 0.84	ND	7/7 (+++)
3	743.6 ± 184.5	728.1 ± 88.0	6.58 ± 0.70	6.46 ± 0.89	ND	3/7 (++)
4	849.0 ± 193.6	704.1 ± 51.9	5.10 ± 1.23	6.08 ± 1.17	1/7 (+)	4/7 (++)

* significantly different from control group of the same age by ANOVA with post hoc Tuckey test.

^a *E. coli* growing on a medium with ciprofloxacin

^b Experimental strain detected in 10⁻⁶ – 10⁻⁵ dilution (+++), 10⁻⁴ – 10⁻³ dilution (++)¹, 10⁻² – 10⁻¹ dilution (+), ND – not detected in any dilution



Journal Pre-proof